

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b>  <b>C12Q 1/68</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/55363</b>  <b>(43) International Publication Date:</b> 21 September 2000 (21.09.00)
<b>(21) International Application Number:</b> PCT/GB00/00807  <b>(22) International Filing Date:</b> 9 March 2000 (09.03.00)  <b>(30) Priority Data:</b> 9905807.5                      12 March 1999 (12.03.99)                      GB  <b>(71) Applicants (for all designated States except US):</b> AMERSHAM PHARMACIA BIOTECH UK LTD [GB/GB]; Amersham Laboratories, White Lion Road, Amersham, Buckinghamshire HP7 9LL (GB). AMERSHAM PHARMACIA BIOTECH INC [US/US]; 800 Centennial Avenue, P.O. Box 1327, Piscataway, NJ 08855-1327 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> THOMAS, Nicholas [GB/GB]; 12 Mapletree Close, Radyr, Cardiff CF4 8RU (GB). WAGGONER, Alan [US/US]; Carnegie Mellon University, 4400 5th Avenue, Pittsburg, PA 15213 (US).  <b>(74) Agents:</b> ROLLINS, Anthony, John et al.; Nycomed Amersham plc, Amersham Laboratories, White Lion Road, Amersham, Buckinghamshire HP7 9LL (GB).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> ANALYSIS OF DIFFERENTIAL GENE EXPRESSION  <b>(57) Abstract</b>  <p>The invention provides methods for detecting the differential expression or presence of two analytes, and more specifically to procedures which provide for rapid and efficient analysis of gene expression in biological systems. In particular, the invention provides a method of detecting and analysing differences between nucleic acids from two sources, which method comprises: a. providing nucleic acids from two sources as labelled probes; b. forming a mixture of the labelled probes with pooled reagents wherein each reagent is a population of beads carrying a polynucleotide target, the target of one reagent being different from the target of another reagent, the beads of one reagent being distinguishable from the beads of another reagent; c. incubating the mixture under conditions to promote specific hybridisation between probes and targets; and, d. analysing beads in the mixture by flow cytometry.</p>		

 J1017 U.S. PTO  
 10/092885  
 03/06/02

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

WO 00/55363

PCT/GB00/00807

### ANALYSIS OF DIFFERENTIAL GENE EXPRESSION

This invention relates to methods for detecting the differential expression or presence of two analytes, and more specifically to procedures which provide for rapid  
5 and efficient analysis of gene expression in biological systems.

Analysis of cellular gene expression is key to gaining understanding of the function of biological organisms and to elucidating the mechanisms which control key cellular events, knowledge of which is crucial to the development of drugs and strategies for treatment of disease states arising from disruption of cellular control  
10 processes.

A wide variety of methods have been developed for analysing gene expression, both at the level of mRNA expression, or by examining the amounts of specific proteins present in cells. Most methods have the same experimental basis in that they examine differential expression; that is they examine the difference in  
15 expression of one or more cellular components between two cells which have been exposed to different conditions. Such studies typically compare expression levels in cells of an untreated population (the control cells) with expression in a separate population of the same cell type (the test cells) that have been subjected to some form of stimulus, for example exposure to a hormone, drug or other chemical.

20 Early methods for the analysis of differential gene expression were predominantly based on analysis of (often anonymous) mRNA bands on electrophoresis gels. Such approaches have largely been displaced by more powerful, reproducible and informative methods based on the use of arrays of nucleic acids in which large numbers of specific sequences are laid down in an ordered pattern on a solid surface and form the  
25 target for hybridisation and capture of labelled mRNA or cDNA from the cells under study. Such arrays have been constructed on a variety of supports, ranging from nylon membranes to glass and silicon wafers. Whatever the support, the essential method of use is the same: firstly known sequences, complementary to cellular mRNAs, either in the form of synthetic oligonucleotides or as PCR products, are laid down on the solid  
30 support in spots at defined locations. These immobilised sequences (targets) are then exposed to sequences (probes) extracted from the cells or tissue under study, where the probes are tagged with some form of label which can be detected in subsequent analysis.

WO 00/55363

PCT/GB00/00807

- 2 -

In early techniques, radioactive labels were used to probe targets on nylon membranes using techniques developed from Southern and Northern blotting methods. However these methods required the hybridisation process to be performed twice, once for the control sample and once for the test sample. They were subsequently overtaken by more elegant procedures based on the use of different coloured fluorophors to label control and test probes and using either PCR products coupled to glass (Schena M. *et al*, (1996), Proc. Natl. Acad. Sci., 93 (20), 10614-10619) or oligonucleotides synthesised directly onto glass using photo-lithographic techniques (Chee M. *et al*, (1996), Science 274 (5287), 610-61) to form what are commonly termed 'micro-arrays'.

In both methods the mRNA sequences extracted from control and test cells or tissues are either labelled directly or are first converted or amplified to yield equivalent cDNA sequences which are subsequently labelled. Once immobilised by hybridisation to complementary target sequences located on the array, the fluorescent labels attached to the probes are detected, either by scanning or by imaging, and quantified to yield data on the amounts of different mRNAs present in the test and control samples. Since sequences from test and control cells are labelled with different fluorophors, both samples can be applied and hybridised simultaneously and the resulting pattern and intensity of hybridised probes determined using detection instrumentation tuned to distinguish between the emission wavelengths of the fluorophors used.

Consequently these two-colour methods allow direct visualisation of differential expression of mRNAs between the two cell populations and are widely used in many fields of life science research to study the control and consequences of gene expression.

However, despite the relative simplicity and elegance of these methods, in operation they suffer from a number of technical difficulties which limit the ease of application and the speed of the techniques to the analysis of gene expression. The construction of target arrays is a time consuming and often expensive process requiring precision equipment; either for construction and alignment of the masks required for light directed oligonucleotide synthesis, or for precise application of nanolitre droplets of liquid for DNA arrays. Arrays based on DNA spots also suffer from artefacts arising from unequal application or drying of the very small volumes used, and often require

WO 00/55363

PCT/GB00/00807

- 3 -

replicate spots to yield accurate data. With either type of array, if the user wishes to modify his investigations to include additional sequences a whole array must be constructed to accommodate the new targets. It would not be atypical to have to make an entirely new array simply to add a single new sequence to a pre-existing library of  
5 several thousand sequences.

The hybridisation process is also subject to a number of problems arising from the geometry of the system and the temperature required for the hybridisation process. To achieve sufficient sensitivity to detect low levels of mRNAs it is necessary to use a high concentration of labelled probe to achieve maximal hybridisation to target  
10 sequences; this requirement and the limited amount of probe material available results in hybridisation reactions being performed in very small volumes. Difficulties therefore arise in ensuring adequate coverage of arrays with microlitre quantities of solution, and the resulting thin films of liquid do not promote good access of the mobile probe sequences to the fixed target sequences; furthermore problems with evaporation are also  
15 common at the temperatures of 40°C to 65°C commonly used for hybridisation.

Finally, the detection and quantification of fluorophor-labelled probes on micro-arrays requires dedicated sophisticated equipment to detect the very low levels of fluorescence present. To achieve the required sensitivity; detection is most commonly achieved using a scanning laser spot to excite fluorophor molecules; this can be a very  
20 slow process requiring up to several hours to complete measurements from a single array.

This invention provides an alternative to micro-array systems for analysis of gene expression. Means are provided for performing analyses using a particle based technique so as to replace ordered 2D arrays with randomly oriented 3D  
25 arrays which can be quickly and easily modified to include new target sequences. The technique provides favourable geometry and kinetics for promoting efficient hybridisation, that can be performed in a standard reaction tube, and that allows measurement of hybridised probe to several thousand target sequences to be accomplished in a few seconds.

30 The invention provides a method of detecting and analysing differences between nucleic acids from two sources, which method comprises:

- a. providing nucleic acids from two sources as labelled probes;
- b. forming a mixture of the labelled probes with pooled reagents wherein

WO 00/55363

PCT/GB00/00807

- 4 -

each reagent is a population of beads carrying a polynucleotide target, the target of one reagent being different from the target of another reagent, the beads of one reagent being distinguishable from the beads of another reagent;

- c. incubating the mixture under conditions to promote specific
- 5 hybridisation between probes and targets; and,
- d. analysing beads in the mixture by flow cytometry.

A polynucleotide target is partly or wholly single-stranded and is capable of specific hybridisation. Oligonucleotides of at least 8 residues are preferred. Preferred are cDNA sequences derived, e.g. by RT-PCR amplification, from cellular

10 mRNA.

The pooled reagents may comprise one bead, or preferably a plurality of beads, of each reagent.

Features of the invention include:

- a) gene expression assays are performed on carrier beads;
- 15 b) individually identifiable beads or populations of beads each carrying a different target sequence are prepared;
- c) selected beads, or populations of beads, are pooled together in suspension to provide a randomly oriented 3D array of particles carrying all sequences of interest for an individual investigation;
- 20 d) mRNAs or cDNAs prepared from control and test cells or tissues are labelled with fluorescent tags to identify their source;
- e) labelled probe species are mixed with the pooled suspension of target carrying beads under conditions which promote specific hybridisation between probes and targets;
- 25 f) the bead mixture is analysed by flow cytometry to simultaneously determine the identity of each bead analysed (and hence the identity of the target sequence carried by the bead) and to quantify the amounts of both control and test probes bound to each bead; and
- g) data is analysed to yield information on the relative and absolute
- 30 abundance of each mRNA in the control and test samples.

The beads of one reagent can be distinguished from the beads of another reagent by a number of different means. Suitable distinguishing means include differences in size, colour or fluorescence or the nature or concentration of markers

WO 00/55363

PCT/GB00/00807

- 5 -

attached to the beads. Beads of one reagent can be distinguished from beads of other reagents using one or more of such means.

For clarity, the invention is now described with reference to the

5 following figures:

Figure 1: Schematic illustration of 2D ordered array and 3D random array.

Figure 2: Flowchart illustrating the principle of the bead based flow cytometry gene expression process.

10 Figure 3: Schematic representation of the bead based flow cytometry gene expression process.

Figure 4: Schematic representation of method for analysis of differential expression of TNF and GAPDH genes in LPS stimulated THP-1 cells.

15 With reference to Figure 1, in a two dimensional ordered array the locations of each target spot immobilised on a planar surface are defined by x,y co-ordinates and hence target sequences are identified by the same co-ordinates. In contrast, in a 3D array formed from particles dispersed in a space defined by dimensions x,y,z, if each particle is individually identifiable by some inherent characteristic, it is not  
20 necessary to use x,y,z locations to specify the identity of each bead and the particles can be randomly distributed throughout the volume as in a suspension of beads in liquid. It follows that if each bead is individually identifiable, then any target sequence previously coupled to that bead is also identifiable. Therefore if a number of differing beads, or discrete populations of beads, are individually prepared where each bead  
25 carries a different target sequence and then selectively pooled, the pooled beads can form a 3D array which can be used for gene expression analysis.

Beads suitable for use in the method of the invention are those which can be readily identified during analysis by flow cytometry: such beads have been previously developed and used for diagnostic assays to measure a wide range of  
30 analytes in blood and other biological fluids by immunoassay. A desire to have a higher throughput in these applications has led to the development of multiplex methods which allow more than one analyte to be measured simultaneously by means of flow cytometry analysis. Multiplexing is achieved by carrying out solid phase linked assays

WO 00/55363

PCT/GB00/00807

- 6 -

using plastic or latex beads as assay substrates. By using a number of discrete bead types which are individually distinguishable from each other, where each bead type carries reagents for one assay, standard flow cytometer instrumentation may be used both to identify the bead type and to measure the assay signal associated with each bead.

- 5 Discrimination between bead populations can be achieved by size (Frengen J. *et al* (1995), Journal of Immunological Methods, Volume 178, p141), by colour or fluorescence (Fulwyier M.J. UK Patent 1,561,042) or by electronic means (Mandecki W. US Patent 5,641,634).

- 10 The general principle of the process of the invention is now described with reference to Figure 2. Selected target cDNA sequences are prepared by standard PCR methods incorporating a means to allow coupling of target sequences to beads. One suitable method would utilise a 5'-biotin on one of the PCR primers, yielding a 5'-biotinylated DNA suitable for coupling to streptavidin-coated beads. Those skilled in  
15 the art will recognise that alternative chemical coupling strategies are available. Such alternative strategies may include, for example, synthesising oligonucleotides having a chemical group such as an amino group at the 5' end thus rendering them suitable for crosslinking to beads which have been modified to have, for example, carboxyl groups on their surface. It will also be appreciated that oligonucleotides synthesised with a  
20 terminal biotin or other coupling group could readily be used in place of PCR generated DNA sequences.

- Once the required number of target cDNAs (cDNA 1 to cDNA n) have been prepared, each target sequence is separately coupled to a corresponding discrete population of beads (Bead 1 to Bead n respectively). Aliquots are then removed from  
25 each population and pooled to form a mixed suspension of beads constituting a randomly orientated 3D array of target sequences. The 3D array is then hybridised with fluorescently labelled probes (RNA or cDNA) prepared from the control and target cells or tissues which have been labelled with two different fluorophors (Fluor A and Fluor B respectively). Following hybridisation the mixed population of beads is analysed by  
30 flow cytometry; as each bead is analysed information from the flow cytometer detectors is used to identify the bead and to measure the amounts of Fluor A (control mRNA) and Fluor B (sample mRNA) bound to the complementary target sequence carried by the bead. These measurements are then used to determine the relative expression of each



WO 00/55363

PCT/GB00/00807

- 7 -

mRNA in the samples.

In a further illustration of the method of the invention reference is made to one embodiment as shown in Figure 3. Control (1) and test (2) probes are prepared and labelled using standard methods and aliquots mixed in a tube (3) containing a mixture of beads (4) carrying the desired target sequences and the tube sealed. Hybridisation of probe and target sequences are promoted by incubating the mixture under conditions of heat, pH and salt concentration which are known to allow the formation of specific nucleic acid hybrids. Following hybridisation, the bead mixture is analysed by flow cytometry using multiple channel fluorescence detection. In the embodiment illustrated, two fluorescence channels are used to identify beads and two further channels are used to measure control and test probe fluorescence. For each bead passing through the flow cytometer this data produces a set of data values that can be represented as 3D plots for control probes (7) and test probes (6). Bead identity is determined by measuring the amounts of two different fluorophors (bead Fluor 1 and bead Fluor 2) incorporated within the bead during manufacture. Plotting the intensities of the two fluorophors on x,y axes (8 & 9) separates the different bead populations used.

The number of possible target sequences that can be measured in a single assay will necessarily be limited by the number of bead populations which it is possible to discriminate in a mixture. With current flow cytometry instrumentation this does not pose a limitation on the utility of the procedure. Typical modern flow cytometry instruments are capable of simultaneously measuring fluorescence at four wavelengths together with other parameters, for example light scattering which is a measure of the size of particles under analysis. In addition, the dynamic range of fluorescence detection is high and fluorescence may be accurately measured over several orders of magnitude. Given this sophistication in measurement it is relatively straightforward to devise schemes which yield a large number of individually distinguishable bead populations to serve as carriers. For example if beads are prepared which contain 2 separate fluorophors, with each fluorophor present in one of 10 levels, then  $10^2 = 100$  bead types are created. By increasing the number of fluorophors, or the levels of each fluor or by introducing other variables, such as bead size, larger numbers of discrete bead types can be produced.

Plotting the intensity of probe fluorophors on the z axis of two different

WO 00/55363

PCT/GB00/00807

- 8 -

plots (10 & 11) shows the amounts of the control (10) and test (11) probes bound to each bead population. This allows the production of a table of the amounts bound to each target sequence in the analysis (14). In the schematic example shown, one mRNA species (12) is expressed at a lower level in the test sample than in the control (13).

- 5 Other differences in expression can be readily identified as differences in the heights of equivalent peaks in the two plots.

In micro-array technique applications DNA or oligonucleotide target sequences are typically applied to a solid surface as discrete areas of dimensions in the range 10-100  $\mu\text{m}$ , with dimensions of 50-100  $\mu\text{m}$  being typical of DNA spots applied as liquid droplets, and smaller areas being used in techniques utilising photo-lithographic oligonucleotide synthesis. To ensure accuracy in measurement of differential expression it is important that the amount of DNA or oligonucleotide present on the solid phase is in excess of complementary sequences in the probe solution such that target sequences do not become limiting leading to distortion of hybridisation results. Consequently it is crucial that in any procedure using flow cytometry for analysis of gene expression using target sequences carried on beads, that the capacity of the system retains the same degree of target:probe excess as used in conventional techniques. Beads used for flow cytometry typically have diameters in the range from 1-10  $\mu\text{m}$  and therefore individually do not have sufficient surface area to substitute for a typical micro-array. However by using several beads to carry each target sequence it is possible to achieve equivalence in target presentation as shown in the following example:

25 spot area      For a 2D array with 1 00  $\mu\text{m}$   $\varnothing$  spots:

$$= \pi r^2$$

$$= 3.14 \times (50)^2$$

$$= 7850 \mu\text{m}^2$$

30 lit area      For 10  $\mu\text{m}$   $\varnothing$  beads in beam (50% surface illuminated):

$$= 2\pi r^2$$

$$= 2 \times 3.14 \times (5)^2$$

$$= 157 \mu\text{m}^2$$

WO 00/55363

PCT/GB00/00807

- 9 -

Bead:Spot equivalence = 7850 / 157  
= 50 beads

5    Bead volume (assuming cubic packing)                    =  $10^3 \mu\text{m}^3$   
      Volume of 50 beads    =  $50,000 \mu\text{m}^3$   
      Assuming 10% v/v suspension                                =  $500,000 \mu\text{m}^3$   
    =  $5 \times 10^5 / 1 \times 10^9 \mu\text{l}$   
    =  $1 \times 10^{-3} \mu\text{l}$   
 10    = 1 nl

Consequently for a complete assay of 1000 targets using 50 beads/target the total volume required for the assay is 1  $\mu\text{l}$ . If desired, larger volumes may be used for convenience in hybridisation or analysis; for example using beads at a concentration  
 15 of 1 % v/v would give a 10  $\mu\text{l}$  total volume.

If desirable, the method of the invention would allow larger numbers of beads to be used with a consequent increase in capacity for binding a greater mass of target sequence spread over the total bead population. This would allow the user if desired to increase the amount of probe bound to the bead populations to increase the  
 20 sensitivity of the process for detecting rare species. Alternatively, it enables an increase in the number of samples which may be analysed simultaneously, for example, to measure expression of a panel of genes simultaneously in a control and more than one test sample, where as described previously each control or test sample is labelled with a different fluorophor. Such increases in assay complexity are not achievable with  
 25 conventional arrays on solid surfaces without reducing sensitivity due to the finite capacity of array spots for binding complementary sequences.

It can be appreciated by the skilled worker that the method of the present invention provides a number of significant advantages over previously described procedures for gene expression analysis which are based on 2D arrays:  
 30 a) the basic components for the bead based assay are readily prepared by coupling solutions of cDNA or oligonucleotides to commercially available beads using standard coupling methods,

WO 00/55363

PCT/GB00/00807

- 10 -

- b) no specialised equipment is required for preparation or analysis, in contrast to the dedicated array production and scanning equipment required for micro-arrays,
- c) the design of investigations can be easily modified with target sequences being added or deleted at will without the requirement to scrap existing materials,
- d) hybridisation is performed in suspension in standard reaction vessels, thereby avoiding problems with evaporation associated with thin films of liquid covering micro-arrays and promoting hybridisation through efficient mixing of probe and target sequences, and
- 10 e) analysis speed is significantly improved: flow cytometers typically analyse beads at rates of 1,000-10,000 beads/second allowing processing of a 100 sequence gene expression analysis in a few seconds.

15 In a further illustration of the method of the invention reference is made to one embodiment as shown in Figure 4.

Figure 4 shows a scheme of a method for the analysis of differential Tumour Necrosis Factor (TNF) expression in THP-1 cells either treated in the presence (test) or absence (control) of bacterial lipopolysaccharide (LPS) (Su S. et al BioTechniques 1997, 22:1107-1113). In this method, variations in abundance of the

20 TNF gene transcripts in control and test samples are simultaneously compared to levels of transcripts from the glyceraldehyde-3-phosphate dehydrogenase (GAPDH) gene in the same samples.

Briefly, human monocyte THP-1 cells at  $5 \times 10^5$  cells/ml are treated in the presence (test sample) and absence (control sample) of  $10 \mu\text{g/ml}$  LPS for 90

25 minutes. Following treatment, both samples are processed separately to isolate the RNA populations present in the control and test samples according to standard procedures set out in Molecular Cloning, A Laboratory Manual, Cold Spring Harbour Laboratory Press 1989, pp 7.3-7.87. Other suitable methods for RNA isolation will be recognised by someone skilled in the art and include the use of commercially available reagents or kits

30 (e.g. RNeasy, Qiagen).

Following RNA isolation, mRNA molecules are converted to cDNA by means of the enzyme, reverse transcriptase, using the standard method set out in Molecular Cloning, A Laboratory Manual, Cold Spring Harbour Laboratory Press 1989.

WO 00/55363

PCT/GB00/00807

- 11 -

pp 8.11-8.13.

The cDNA molecules obtained from each sample are then used in separate multiplex PCR reactions using standard conditions (Molecular Cloning, A Laboratory Manual, Cold Spring Harbour Laboratory Press 1989 pp 14.5-14.20) and primer pairs which amplify GAPDH and TNF (Su S. et al BioTechniques 1997, 22:1107-1113). One primer of each primer pair is modified at the 5' end with a biotin molecule to aid in strand separation in subsequent steps in the process. This is readily achieved using standard techniques during oligonucleotide synthesis with modified phosphoramidites, for example : 5'-Biotin Phosphoramidite (Glen Research).

Fluorescently labelled nucleotides are incorporated into the PCR reactions in order to label the amplified PCR products such that their origin from the test or control RNA populations may be established. Thus, the PCR reaction mix containing cDNAs prepared from the control cells includes Cy3<sup>TM</sup>-dCTP and the reaction mix containing cDNAs prepared from test cells includes Cy5<sup>TM</sup>-dCTP (Cy3<sup>TM</sup>-dCTP and Cy5<sup>TM</sup>-dCTP are obtained from Amersham Pharmacia Biotech). These reactions yield amplified levels of fluorescently labelled cDNA populations wherein the relative abundance of individual cDNA species corresponds to the abundance of their respective parent mRNA species in the RNA populations isolated from the cell or tissue samples and where the origin of any given molecule from the test or control sample populations can be ascribed from the fluorescent label carried by the molecule.

The PCR products from the control (Cy3-labelled) and test (Cy-5 labelled) reactions are now mixed together prior to further processing.

Double stranded cDNA products of the PCR reactions are converted to single stranded cDNA molecules prior to further analysis by binding the PCR products via the biotin at the 5' end of each PCR product to streptavidin-coated magnetic beads (MagneSphere, Promega). Once bound, the double stranded PCR products are denatured by addition of 0.2 volume of 2M NaOH, incubating for 10 minutes at room temperature in order to release the non-biotinylated strands from the beads. The beads are separated from the solution by attraction to a magnet and the clarified solution, which contains the single stranded labelled PCR products, is removed and made neutral by addition of 1 volume of 0.4M HCl. Other methods for separating double-stranded DNA molecules are known to those skilled in the art.

Populations of beads or particles suitable for performing analysis of

WO 00/55363

PCT/GB00/00807

- 12 -

differential gene expression are prepared using beads containing fluorescent dyes (SPHERO™, Spherotech Inc.). Yellow fluorescent streptavidin-coated beads are coated with the 5'-biotin primer for GAPDH and blue fluorescent streptavidin-coated beads are coated with the 5'-biotin primer for TNF (using the same 5'-biotin primers as were used earlier). Thus, these populations of beads have two distinguishing characteristics i.e. each bead population is distinguishable from all other bead populations by its fluorescence characteristics and each bead population is capable of recognising a single cDNA species (either TNF or GAPDH).

The separately prepared beads are mixed together, further mixed with the Cy 3 and Cy 5-labelled single stranded cDNA products prepared previously and incubated in a hybridisation buffer (0.1M Tris.HCl pH7.4, 750mM NaCl) at 45° for 2 hours prior to analysis of the bead populations by flow cytometry. cDNA products from the mixed population will bind specifically to their complementary capture sequences carried on beads such that each bead will ultimately be decorated with a mixture of labelled cDNAs of a single species arising from the test and control populations, and where the relative abundance of the labels reflects the relative abundance of single species in the original test and control populations.

The relative abundance of GAPDH transcripts in the population of molecules derived from control and test samples is compared by detecting the fluorescence of yellow beads at 460nm/ 480nm (excitation/emission); and, within the population of yellow beads, bound Cy3 labelled transcripts are detected at 550nm/570nm (excitation/emission) and bound Cy5 labelled transcripts are detected at 650nm/670nm (excitation/emission). The relative abundance of TNF transcripts in the molecules derived from test and control samples is determined by detecting the fluorescence of blue beads at 650nm/710nm (excitation/emission); and further detecting bound Cy3 labelled transcripts at 550nm/570nm (excitation/emission) and bound Cy5 labelled transcripts at 650nm/670nm (excitation/emission). Thus bead fluorescence is used to assign the identity of the gene associated with each bead and label fluorescence is used to determine the relative abundance of cDNAs arising from the test and control samples which are attached to the bead, the latter yielding information on the differential expression of that gene under the conditions used in treatment of the original cell or tissue sample.

WO 00/55363

PCT/GB00/00807

- 13 -

CLAIMS

1. A method of detecting and analysing differences between nucleic acids  
5 from two sources, which method comprises:
  - a. providing nucleic acids from two sources as labelled probes;
  - b. forming a mixture of the labelled probes with pooled reagents wherein  
each reagent is a population of beads carrying a polynucleotide target, the target of one  
reagent being different from the target of another reagent, the beads of one reagent  
10 being distinguishable from the beads of another reagent;
  - c. incubating the mixture under conditions to promote specific  
hybridisation between probes and targets; and,
  - d. analysing beads in the mixture by flow cytometry.
- 15 2. The method of claim 1 wherein the nucleic acids from two sources are  
mRNA or cDNA from cells or tissues.
3. The method of claim 1 or claim 2 wherein the polynucleotide targets are  
cDNA derived from cellular mRNA.
- 20 4. The method of any one of claims 1 to 3 wherein the polynucleotide  
targets are PCR amplimers.
5. The method of any one of claims 1 to 4 wherein the polynucleotide  
25 targets carry terminal biotin groups through which they are attached to  
streptavidin-coated beads.
6. The method of any one of claims 1 to 5 wherein the polynucleotide  
targets are single-stranded nucleic acids.
- 30 7. The method of any one of claims 1 to 6 wherein the labelled probes are  
single-stranded nucleic acids.

WO 00/55363

PCT/GB00/00807

- 14 -



8. The method of any one of claims 1 to 7 wherein beads of one reagent are distinguishable from beads of another reagent by size.
9. The method of any one of claims 1 to 8 wherein beads of one reagent are distinguishable from beads of another reagent by the nature of the markers attached to the beads.
10. The method of any one of claims 1 to 9 wherein beads of one reagent are distinguishable from beads of another reagent by the concentration of markers attached to the beads.
11. The method of any one of claims 1 to 7 wherein beads of one reagent are distinguishable from beads of another reagent by size and/or by the nature and/or the concentration of markers attached to the beads.
12. The method of any one of claims 8 to 11 wherein fluorescent markers are attached to the beads.
13. The method of claim 1 or claim 2 wherein each probe is labelled with a fluorescent tag to indicate its source.
14. The method of any one of claims 1 to 13 wherein analysis by flow cytometry is performed to identify each bead and to quantify the probes bound thereto.
15. The method of any one of claims 1 to 14 wherein data obtained by flow cytometry is analysed to yield information about the relative and/or absolute abundances of individual sequences of the nucleic acids from the two sources.



WO 00/55363

1/4


PCT/GB00/00807

**Figure 1.****2D Ordered Array****3D Random Array**

WO 00/55363

2/4


PCT/GB00/00807

**Figure 2**

WO 00/55363

3/4

PCT/GB00/00807

**Figure 3.**

WO 00/55363

4/4

PCT/GB00/00807

**Figure 4.** Analysis of differential expression of TNF & GAPDH genes in LPS stimulated THP-1 cells

